

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Hybanthus, Ipomoea, Cordia, Russelia, Tetramerium, Isertia, Liabum (2). IV. "Diagnoses of new spermatophytes from Mexico," by M. L. Fernald, new species being described under Carex, Alnus (2), Heliotropium, Salvia (9), Castilleja, Ruellia.—J. M. C.

Bicentennary of Linné.—In connection with the Linnean celebration at the University of Upsala, a series of eight publications has been issued. In general the volumes contain reprints of some of the most interesting minor papers of Linné, which thus become accessible to a far greater number of readers. For example, the "Invitation du recteur pour assister aux fêtes" is a paper of 107 pages, consisting chiefly of a reprint of Linné's "Cultur der Pflanzen." The announcements in reference to the conferring of doctor's degrees in philosophy, medicine, law, and theology are four volumes of reprints. There is also a special publication, "Linné och Vaxtodlingen," edited by Swederus. The first (pp. 341) of four volumes containing the correspondence of Linné is also included. The most elaborate member of the series is the "Linnéporträtt," prefaced by a colored portrait of Linné, and containing reproductions of numerous other portraits, busts, medals, etc., as well as a description of the 515 portraits (paintings, medals, etc.) in the collection of the University. The University of Upsala has certainly spared no labor and expense in doing honor to her illustrious professor.—J. M. C

Genera Siphonogamarum.—The tenth fascicle of Dalla Torre and Harms's<sup>6</sup> list of the genera of seed-plants continues the general alphabetical list of names, the last entry being Macrocarpium.—J. M. C.

## NOTES FOR STUDENTS

Fungi in termite nests.—Petch gives an account of the fungi found in certain termite nests in Ceylon, which grow from the combs found in the chambers. The fungus flora of the combs in their normal state seems to be limited to few species which occur almost pure. The only form on the normal comb is a hyphomycete which was not determined, but from the descriptions seems to be like Sterigmatocystis. This fungus seems to be endemic in the nests, according to the author not being found outside them.

When the combs grow old they give rise to two forms of agarics, which, however, the author regards as one species. Both have been described under several names from material sent to Europe. These agarics arise from combs at a considerable depth below the surface, so that their rootlike stalks attain an average length of 30<sup>cm</sup>. The lower part of the stalk is black, while the upper portion is white. The first form is marked by the absence of an annulus and by the peculiar fact that only a single plant develops from each comb. Although a large number start, only one pushes its way to the surface of the ground. All the other rudiments fail entirely to develop, so that it is not possible to find specimens which have

<sup>&</sup>lt;sup>6</sup> Dalla Torre, C. G. de, and Harms, H., Genera Siphonogamarum ad systema Englerianum conscripta. Fasc. 10. pp. 721–800. Leipzig: Wilhelm Engelmann. 1907. *M* 6.